

PRODUCT INFORMATION

Target VSIG4

Synonyms CRIg;Z39IG

Recombinant Human VSIG4 with C-terminal **Description**

human Fc tag

Delivery In Stock **Uniprot ID** Q9Y279 **Expression Host HEK293**

Tag C-Human Fc Tag

Molecular

Storage & Shipping

Background

VSIG4(Arg20-Pro283) hFc(Glu99-Ala330) Characterization

The protein has a predicted molecular mass of **Molecular Weight**

55.3 kDa after removal of the signal peptide. The apparent molecular mass of VSIG4-hFc is

approximately 55-70 kDa due to glycosylation. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a v-set and immunoglobulin-domain containing protein that is structurally related to the B7 family of immune regulatory proteins. The encoded protein may be a negative regulator of T-cell responses. This protein is also a

receptor for the complement component 3 fragments C3b and iC3b. Alternate splicing results in multiple transcript variants. [provided

by RefSeq, May 2010]

Usage Research use only

Conjugate Unconjugated














Figure 1. Human VSIG4 Protein, hFc Tag on SDS-PAGE under reducing condition.

Human VSIG4,hFc Tagged protein ELISA

0.2 µg of Human VSIG4, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human VSIG4 Protein, hFc Tag(PME100855) can bind Anti-VSIG4 antibody(DMC268), IgG1 Chimeric mAb in a linear range of 3.20–80 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com

