

## **PRODUCT INFORMATION**

**Clone ID** DM4 **BCMA Target** TNFRSF17 **Synonyms Host Species** Rabbit

PE-conjugated Anti-BCMA bispecific antibody(DM4) Description

**Delivery** 3-4 weeks **Uniprot ID** Q02223 **IgG** type Rabbit scfv Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended Flow Cyt 1:100 **Dilutions** 

Formulation &

**Background** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Liquid PBS with 0.05% Proclin300, 1% BSA Reconstitution

Storage & Shipping Store at 2°C-8°C for 6 months

BiTE bispecific antibody is an Engineered fusion protein constructed from two single-chain variable fragments (scFvs) of different monoclonal antibodies. One of the scFvs will be constructed from an anti-CD3 monoclonal antibody; and the other scFv fragment which linked by a linker region will be made from an anti-Tumor cell specific monoclonal antibody. The B-cell maturation protein (BCMA or BCM) is a member of the TNF-receptor superfamily. This

receptor is preferentially expressed in mature B lymphocytes; and may be important for B cell development and autoimmune response. This receptor has been shown to specifically bind to the tumor necrosis factor (ligand) superfamily; member 13b (TNFSF13B:TALL-1:BAFF); and to lead to NE kappaP and MARKS:INK activation. The lead to NF-kappaB and MAPK8:JNK activation. This receptor also binds to various TRAF family

members; and thus may transduce signals for cell survival and proliferation. [provided by RefSeq; Jul

> Email: info@dimabio.com Website: www.dimabio.com

20081

**Usage** Research use only Conjugate PE-conjugated

