

## **PRODUCT INFORMATION**

Clone ID DM102 **CD40 Target** 

**Synonyms** CD40; Bp50; CDW40; MGC9013; TNFRSF5; p50

**Host Species** 

PE-conjugated Anti-CD40 antibody(DM102); Description

Rabbit mAb 3-4 weeks

**Delivery Uniprot ID** P25942 Rabbit IgG IgG type Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended Flow Cyt 1:100 **Dilutions** 

Formulation &

**Background** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Liquid PBS with 0.05% Proclin300, 1% BSA Reconstitution

Storage & Shipping Store at 2°C-8°C for 6 months

This gene is a member of the TNF-receptor superfamily. The encoded protein is a receptor on antigen-presenting cells of the immune system and is essential for mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching; memory B cell development; and germinal center formation. AT-hook transcription factor AKNA is reported to coordinately regulate the expression of this receptor and its ligand; which may be important for homotypic cell interactions. Adaptor protein TNFR2 interacts with this receptor and serves as a mediator of the signal transduction.

The interaction of this receptor and its ligand is found to be necessary for amyloid-beta-induced microglial activation; and thus is thought to be an early event in Alzheimer disease pathogenesis. Mutations affecting this gene are the cause of autosomal recessive hyper-IgM immunodeficiency type 3 (HIGM3). Multiple alternatively spliced transcript variants of this gene encoding distinct

> Email: info@dimabio.com Website: www.dimabio.com

isoform's have been reported.

Usage Research use only Conjugate PE-conjugated

