

**PRODUCT INFORMATION**

<b>Clone ID</b>	5B10
<b>Target</b>	CD93
<b>Synonyms</b>	C1qR(P);C1QR1;C1qRP;CDw93;dj737E23.1;EC5M3;MXRA4
<b>Host Species</b>	Rabbit
<b>Description</b>	PE-conjugated Anti-CD93 antibody(5B10), IgG1 Chimeric mAb
<b>Delivery</b>	3-4 weeks
<b>Uniprot ID</b>	Q9NPY3
<b>IgG type</b>	Rabbit/Human Fc chimeric IgG1
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	Flow Cyt
<b>Recommended Dilutions</b>	Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Liquid□PBS with 0.05% Proclin300, 1% BSA
<b>Storage &amp; Shipping</b>	Store at 2°C-8°C for 6 months
<b>Background</b>	The protein encoded by this gene is a cell-surface glycoprotein and type I membrane protein that was originally identified as a myeloid cell-specific marker. The encoded protein was once thought to be a receptor for C1q, but now is thought to instead be involved in intercellular adhesion and in the clearance of apoptotic cells. The intracellular cytoplasmic tail of this protein has been found to interact with moesin, a protein known to play a role in linking transmembrane proteins to the cytoskeleton and in the remodeling of the cytoskeleton. [provided by RefSeq, Jul 2008]
<b>Usage</b>	Research use only
<b>Conjugate</b>	PE-conjugated

