Cat. No. DMC100366P



PRODUCT INFORMATION

Clone ID **DMC366 Target CFB**

AHUS4; ARMD14; BF; BFD; CFAB; CFBD; FB; **Synonyms**

FBI12; GBG; H2-Bf; PBF2

Host Species

PE-conjugated Anti-CFB antibody(DMC366); IgG1 **Description**

Chimeric mAb

Delivery 3-4 weeks **Uniprot ID** P00751

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended

Background

Flow Cyt 1:100 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Formulation & Liquid PBS with 0.05% Proclin300, 1% BSA Reconstitution

Storage & Shipping Store at 2°C-8°C for 6 months

> This gene encodes complement factor B; a component of the alternative pathway of complement activation. Factor B circulates in the blood as a single chain polypeptide. Upon

activation of the alternative pathway; it is cleaved by complement factor D yielding the noncatalytic chain Ba and the catalytic subunit Bb. The active subunit Bb is a serine protease which associates with C3b to form the alternative pathway C3 convertase. Bb is involved in the proliferation of

preactivated B lymphocytes; while Ba inhibits their proliferation. This gene localizes to the major histocompatibility complex (MHC) class III region on coronnosmed is regulated in the several genes involved in regulation of the immune reaction. Polymorphisms in this gene are associated with a reduced risk of age-related macular degeneration. The polyadenylation site of this gene is 421 bp from the 5' end of the gene

> Email: info@dimabio.com Website: www.dimabio.com

for complement component 2.

Usage Research use only Conjugate PE-conjugated

