

## **PRODUCT INFORMATION**

**Clone ID** DM165 CLEC12A **Target** 

**Synonyms** CLEC12A;MICL;CLL-1;CLL1;DCAL2;DCAL-2;CD371

**Host Species** 

PE-conjugated Anti-CLEC12A antibody(DM165); Rabbit mAb Description

**Delivery** 3-4 weeks **Uniprot ID** Q5QGZ9 IgG type Rabbit IgG Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended Flow Cyt 1:100 **Dilutions** 

Formulation &

**Background** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Liquid PBS with 0.05% Proclin300, 1% BSA Reconstitution

Storage & Shipping Store at 2°C-8°C for 6 months

> This gene encodes a member of the C-type lectin:C-type lectin-like domain (CTL:CTLD) superfamily. Members of this family share a common protein fold and have diverse functions; such as cell adhesion; cell-cell signaling;

glycoprotein turnover; and roles in inflammation and immune response. The protein encoded by this gene is a negative regulator of granulocyte and monocyte function. Several alternatively and transcript variants of this gene base have been spliced transcript variants of this gene have been described; but the full-length nature of some of these variants has not been determined. This gene is closely linked to other CTL:CTLD

> Email: info@dimabio.com Website: www.dimabio.com

superfamily members in the natural killer gene complex region on chromosome 12p13.

**Usage** Research use only

PE-conjugated Conjugate

