

PRODUCT INFORMATION

| | |
|---|---|
| Clone ID | DMC388 |
| Target | EDA |
| Synonyms | ED1; EDA2 |
| Host Species | Rabbit |
| Description | PE-conjugated Anti-EDA antibody(DMC388); IgG1 Chimeric mAb |
| Delivery | 3-4 weeks |
| Uniprot ID | Q92838 |
| IgG type | Rabbit/Human Fc chimeric IgG1 |
| Clonality | Monoclonal |
| Reactivity | Human |
| Applications | Flow Cyt |
| Recommended Dilutions | Flow Cyt 1:100 |
| Purification | Purified from cell culture supernatant by affinity chromatography |
| Formulation & Reconstitution | Liquid□PBS with 0.05% Proclin300, 1% BSA |
| Storage & Shipping | Store at 2°C-8°C for 6 months |
| Background | The protein encoded by this gene is a type II membrane protein that can be cleaved by furin to produce a secreted form. The encoded protein; which belongs to the tumor necrosis factor family; acts as a homotrimer and may be involved in cell-cell signaling during the development of ectodermal organs. Defects in this gene are a cause of ectodermal dysplasia; anhidrotic; which is also known as X-linked hypohidrotic ectodermal dysplasia. Several transcript variants encoding many different isoforms have been found for this gene. |
| Usage | Research use only |
| Conjugate | PE-conjugated |

