

PRODUCT INFORMATION

| | |
|---|--|
| Clone ID | DMC487 |
| Target | FZD10 |
| Synonyms | CD350; FZ-10; Fz10; FzE7; hFz10 |
| Host Species | Rabbit |
| Description | PE-conjugated Anti-FZD10 antibody(DMC487); IgG1 Chimeric mAb |
| Delivery | 3-4 weeks |
| Uniprot ID | Q9ULW2 |
| IgG type | Rabbit/Human Fc chimeric IgG1 |
| Clonality | Monoclonal |
| Reactivity | Human |
| Applications | Flow Cyt |
| Recommended Dilutions | Flow Cyt 1:100 |
| Purification | Purified from cell culture supernatant by affinity chromatography |
| Formulation & Reconstitution | Liquid□PBS with 0.05% Proclin300, 1% BSA |
| Storage & Shipping | Store at 2°C-8°C for 6 months |
| Background | This gene is a member of the frizzled gene family. Members of this family encode 7-transmembrane domain proteins that are receptors for the Wingless type MMTV integration site family of signaling proteins. Most frizzled receptors are coupled to the beta-catenin canonical signaling pathway. Using array analysis; expression of this intronless gene is significantly up-regulated in two cases of primary colon cancer. [provided by RefSeq; Jul 2008] |
| Usage | Research use only |
| Conjugate | PE-conjugated |

