

PRODUCT INFORMATION

Clone ID **DMC498 GDNF Target**

Synonyms ATF; ATF1; ATF2; HFB1-GDNF; HSCR3

Host Species

PE-conjugated Anti-GDNF antibody(DMC498); IgG1 Chimeric mAb Description

Delivery 3-4 weeks **Uniprot ID** P39905

Rabbit/Human Fc chimeric IgG1 IgG type

Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended

Background

Flow Cyt 1:100 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Formulation & Reconstitution

Liquid

□PBS with 0.05% Proclin300, 1% BSA

Storage & Shipping Store at 2°C-8°C for 6 months

> beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family

> This gene encodes a secreted ligand of the TGF-

transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer. The recombinant form of this protein; a highly conserved neurotrophic factor; was shown to promote the survival and differentiation of dopaminergic neurons in culture; and was able to

prevent apoptosis of motor neurons induced by axotomy. This protein is a ligand for the product of the RET (rearranged during transfection)
protooncogene. Mutations in this gene may be
associated with Hirschsprung disease and Tourette syndrome. This gene encodes multiple protein isoforms that may undergo similar proteolytic processing. [provided by RefSeq; Aug 2016]

> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only Conjugate PE-conjugated

