

PRODUCT INFORMATION

| | |
|---|--|
| Clone ID | DM126 |
| Target | IL17RA |
| Synonyms | CD217;CDw217;IL-17RA;IL17R;CANDF5;hIL-17R |
| Host Species | Rabbit |
| Description | PE-conjugated Anti-IL17RA antibody(DM126); Rabbit mAb |
| Delivery | 3-4 weeks |
| Uniprot ID | Q96F46 |
| IgG type | Rabbit IgG |
| Clonality | Monoclonal |
| Reactivity | Human |
| Applications | Flow Cyt |
| Recommended Dilutions | Flow Cyt 1:100 |
| Purification | Purified from cell culture supernatant by affinity chromatography |
| Formulation & Reconstitution | Liquid□PBS with 0.05% Proclin300, 1% BSA |
| Storage & Shipping | Store at 2°C-8°C for 6 months |
| Background | <p>Interleukin 17A (IL-17A) is a proinflammatory cytokine secreted by activated T-lymphocytes. It is a potent inducer of the maturation of CD34-positive hematopoietic precursors into neutrophils. The transmembrane protein encoded by this gene (interleukin 17A receptor; IL17RA) is a ubiquitous type I membrane glycoprotein that binds with low affinity to interleukin 17A. Interleukin 17A and its receptor play a pathogenic role in many inflammatory and autoimmune diseases such as rheumatoid arthritis. Like other cytokine receptors; this receptor likely has a multimeric structure. Alternative splicing results in multiple transcript variants encoding different isoforms.</p> |
| Usage | Research use only |
| Conjugate | PE-conjugated |

