

**PRODUCT INFORMATION**

<b>Clone ID</b>	DMC225
<b>Target</b>	IL21R
<b>Synonyms</b>	CD360; IMD56; NILR
<b>Host Species</b>	Rabbit
<b>Description</b>	PE-conjugated Anti-IL21R antibody(DMC225); IgG1 Chimeric mAb
<b>Delivery</b>	3-4 weeks
<b>Uniprot ID</b>	Q9HBE5
<b>IgG type</b>	Rabbit/Human Fc chimeric IgG1
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	Flow Cyt
<b>Recommended Dilutions</b>	Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Liquid□PBS with 0.05% Proclin300, 1% BSA
<b>Storage &amp; Shipping</b>	Store at 2°C-8°C for 6 months
<b>Background</b>	The protein encoded by this gene is a cytokine receptor for interleukin 21 (IL21). It belongs to the type I cytokine receptors; and has been shown to form a heterodimeric receptor complex with the common gamma-chain; a receptor subunit also shared by the receptors for interleukin 2; 4; 7; 9; and 15. This receptor transduces the growth promoting signal of IL21; and is important for the proliferation and differentiation of T cells; B cells; and natural killer (NK) cells. The ligand binding of this receptor leads to the activation of multiple downstream signaling molecules; including JAK1; JAK3; STAT1; and STAT3. Knockout studies of a similar gene in mouse suggest a role for this gene in regulating immunoglobulin production. Three alternatively spliced transcript variants have been described.
<b>Usage</b>	Research use only
<b>Conjugate</b>	PE-conjugated

