

**PRODUCT INFORMATION**

<b>Target</b>	S1 protein NTD
<b>Synonyms</b>	S1 protein NTD;Spike protein S1 NTD;BetaCoV S1-NTD;COVID-19
<b>Description</b>	Recombinant SARS-CoV-2 (2019-nCoV) S1 protein NTD with C-terminal mouse Fc tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P0DTC2
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-Mouse Fc Tag
<b>Molecular Characterization</b>	S1 protein NTD(Ser13-Leu303) mFc(Pro99-Lys330)
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 59.1 kDa after removal of the signal peptide.The apparent molecular mass of S1-NTD-mFc is approximately 70-100 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as Covid19 (2019 Novel Coronavirus) is a virus that causes illnesses ranging from the common cold to severe diseases. The spike protein is a type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which accounts for recognizing the cell surface receptor, ACE2. S2 contains basic elements needed for the membrane fusion. Recent publications indicate that S1-RBD domain can induce virus neutralizing-antibody and T cell response.
<b>Usage</b>	Research use only






Figure 1. SARS-CoV-2 (2019-nCoV) S1 protein NTD, mFc Tag on SDS-PAGE under reducing condition.

## SARS-CoV-2 (2019-nCoV) S1 NTD, mFc Tagged protein ELISA




Figure 2. ELISA plate pre-coated by 2  $\mu$ g/ml (100  $\mu$ l/well) Human AXL, His tagged protein (PME100070) can bind Human NTD, mFc Tagged protein (PME100492) in a linear range of 0.006-0.16  $\mu$ g/ml.

